

# Prevenar 13<sup>®</sup> Supplementary Dose Tables

valid 1<sup>st</sup> October 2011 to 30<sup>th</sup> September 2012 only

## 1. Standard schedule

Current age of child	Vaccination history	Supplementary dose
12 to 35 months (inclusive)	Previously vaccinated with Prevenar <sup>®</sup>	Give Prevenar 13 <sup>®</sup> at least 8 weeks after last dose of Prevenar <sup>®</sup>

## 2. Aboriginal and Torres Strait Islander (Indigenous) children

### (a) Child has no history of Pneumovax 23<sup>®</sup> vaccination

Current age of child	Vaccination history	Supplementary dose	Pneumovax 23 <sup>®</sup>
12 to 35 months (inclusive)	Previously vaccinated with Prevenar <sup>®</sup>	Give Prevenar 13 <sup>®</sup> at least 8 weeks after last dose of Prevenar <sup>®</sup>	Not required

### (b) Child has already received Pneumovax 23<sup>®</sup> vaccination

Current age of child	Vaccination history	Supplementary dose
12 to 35 months (inclusive)	Previously vaccinated with Prevenar <sup>®</sup> AND Pneumovax 23 <sup>®</sup>	Give Prevenar 13 <sup>®</sup> at least 8 weeks after Pneumovax 23 <sup>®</sup> was given

## 3. Medical at Risk children\* (including children who are both Indigenous AND Medical at Risk)

Current age of child	Vaccination history	Supplementary dose	Pneumovax 23 <sup>®</sup>
12 to 47 months (inclusive)	Previously vaccinated with Prevenar <sup>®</sup>	Give Prevenar 13 <sup>®</sup> at least 8 weeks after last dose of Prevenar <sup>®</sup>	Give at 4-5 years of age <b>OR</b> 8 weeks after last dose of Prevenar 13 <sup>®</sup> (whichever is later)

\*Refer to The Australian Immunisation Handbook (page 246) for details of medical conditions predisposing children to IPD.