


MULTIPLE INJECTION SITES FOR VACCINATION

BEFORE YOU BEGIN

- Is an anaphylaxis kit available?
- Does this child require any catch-up vaccinations?
- Have the vaccines be stored between +2°C to +8°C?
- Are you giving any vaccines that require reconstitution?
- Are there any contraindications or precautions for this child?
- Has this child received the hepatitis B birth dose (given within 7 days of birth)?
- Have you discussed the risks/benefits of vaccination with the parent/carer and documented consent?
- Refer to the Australian Immunisation Handbook, 9th edition 2008 for further information


The child should remain in the practice for a minimum of 15 minutes after vaccination for post-vaccination monitoring.

2, 4, 6 MONTHS




- DTPa / Hib / Hep B / IPV (IM)
- Pneumococcal 13vPCV (IM)
- Also give:
Rotavirus Vaccine (Given Orally)

12 MONTHS



- Hib (IM)
- Measles / Mumps / Rubella (IM)
- Meningococcal C (IM)

18 MONTHS



- Varicella (SC)

4 YEARS OLD



- DTPa / IPV (IM)
- Measles / Mumps / Rubella (IM)

INTRAMUSCULAR (IM) ANTEROLATERAL THIGH OR DELTOID

Angle of needle 90° to the skin
23 or 25 gauge needle
25 mm in length
(16mm may be used for small infants)

SUBCUTANEOUS (SC)

Angle of needle 45° to the skin
25 or 26 gauge needle
16mm in length

